

Arithmetical properties of Multiple Ramanujan sums

Yoshinori YAMASAKI

February 2, 2008

Abstract

In the present paper, we introduce a multiple Ramanujan sum for arithmetic functions, which gives a multivariable extension of the generalized Ramanujan sum studied by D. R. Anderson and T. M. Apostol. We then find fundamental arithmetic properties of the multiple Ramanujan sum and study several types of Dirichlet series involving the multiple Ramanujan sum. As an application, we evaluate higher-dimensional determinants of higher-dimensional matrices, the entries of which are given by values of the multiple Ramanujan sum.

2000 Mathematics Subject Classification : *Primary* 11A25; *Secondary* 11C20.

Key words and phrases : *Ramanujan sum, divisor function, Dirichlet convolution, Dirichlet series, Smith determinant, hyperdeterminant.*

1 Introduction

In 1918, S. Ramanujan [17] studied the sum

$$c(k, n) := \sum_{\substack{l \pmod{k} \\ \gcd(l, k)=1}} e(k, nl) = \sum_{d \mid \gcd(k, n)} \mu\left(\frac{k}{d}\right) d, \quad (1.1)$$

where, in the first sum, l runs over a reduced residue system modulo k with $\gcd(l, k) = 1$, $e(r, n) := \exp(2\pi\sqrt{-1}n/r)$ and μ is the Möbius function. The sum $c(k, n)$ is called the Ramanujan sum (or the Ramanujan trigonometric sum) and is widely investigated in connection with, for example, even arithmetic functions [8, 9] and cyclotomic polynomials [14, 15]. See [13] for the arithmetic theory of the Ramanujan sum. Among several generalizations and variations of $c(k, n)$, D. R. Anderson and T. M. Apostol [1] (see also [3]) considered the sum

$$S_{f,g}(k, n) := \sum_{d \mid \gcd(k, n)} f\left(\frac{k}{d}\right) g(d), \quad (1.2)$$

where f and g are arithmetic functions. Clearly, $S_{f,g}(k, n)$ extends the right-most expression in formula (1.1) and hence gives a generalization of the Ramanujan sum.

Motivated by the study of the above generalized Ramanujan sum, in the present paper, we examine the following type of multiple sum for arithmetic functions $f_1, \dots, f_{m+1}$:

$$S_{f_1, \dots, f_{m+1}}(n_1, \dots, n_{m+1}) := \sum_{\substack{d_j \mid \gcd(n_1, \dots, n_{j+1}) \\ (j=1, \dots, m)}} f_1\left(\frac{n_1}{d_1}\right) f_2\left(\frac{d_1}{d_2}\right) \cdots f_m\left(\frac{d_{m-1}}{d_m}\right) f_{m+1}(d_m),$$

where $\gcd(n_1, \dots, n_{j+1})$ is the greatest common divisor of $n_1, \dots, n_{j+1}$. We call this a *multiple Ramanujan sum* for $f_1, \dots, f_{m+1}$. Notice that the above expression gives the generalized Ramanujan sum (1.2) when $m = 1$ and, moreover, the Dirichlet convolution $f_1 * \dots * f_{m+1}$ of $f_1, \dots, f_{m+1}$ in the “diagonal case”; $n_1 = \dots = n_{m+1}$.

The present paper is organized as follows. In Section 2, we introduce a multiple Ramanujan sum $S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}$ with positive integer parameters $\gamma_1, \dots, \gamma_m$ so that $S_{f_1, \dots, f_{m+1}} = S_{f_1, \dots, f_{m+1}}^{(1, \dots, 1)}$ and study its fundamental properties as a multivariable arithmetic function, such as the degeneracies and the multiplicativity (see [21] for the theory of multivariable arithmetic functions). Then, since $S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}$ belongs to the class of even arithmetic functions (mod n_1) as a function of $n_2, \dots, n_{m+1}$ in the sense of Cohen [9], we calculate its finite Fourier expansion, which any even arithmetic function possesses. This expression is important with respect to Section 4. In Section 3, we study several types of Dirichlet series having coefficients that are given by $S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}$. We treat not only single-variable Dirichlet series, but also multivariable Dirichlet series. For instance, an analogue of the formula of J. M. Borwein and K. K. Choi [4], which contains the classical Ramanujan formula concerning the divisor function σ_a , is obtained. Section 4 is devoted to a higher-dimensional generalization of the so-called Smith determinant [19]. We evaluate higher-dimensional determinants (hyperdeterminants) of higher-dimensional matrices (hypermatrices), the entries of which are given by values of $S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}$. In fact, we derive a hyperdeterminant formula for even multivariable arithmetic functions. This includes the results of P. J. McCarthy [12] and K. Bourque and S. Ligh [5] for the 2-dimensional case, that is, the usual determinant case, and partially of P. Haukkanen [10], for the higher-dimensional case.

We use the following notations in the present paper. The set of natural numbers, the ring of rational integers, the field of real numbers and the field of complex numbers are denoted respectively as $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{R}$, and $\mathbb{C}$. For $n_1, \dots, n_k \in \mathbb{N}$, $\gcd(n_1, \dots, n_k)$ (resp. $\text{lcm}(n_1, \dots, n_k)$) represents for the greatest common divisor (resp. least common multiple) of $n_1, \dots, n_k$. For $x \in \mathbb{R}$, $[x]$ is the greatest integer not exceeding x . We denote the Möbius function as $\mu(n)$, the Euler totient function as $\varphi(n)$, the power function as $\delta^x(n) := n^x$ for $x \in \mathbb{C}$, and the identity element in the ring of arithmetic functions with respect to the Dirichlet convolution $*$ as $\varepsilon(n) := [\frac{1}{n}] = 1$ if $n = 1$, and 0 otherwise. Note that $\delta^0 * \mu = \varepsilon$. Throughout the present paper, we consider a product (resp. a sum) over an empty set to always be equal to 1 (resp. 0).

2 A multiple Ramanujan sum

2.1 Preliminary: γ -convolutions

Let $\mathcal{A}$ be the set of all complex-valued arithmetic functions $f : \mathbb{N} \rightarrow \mathbb{C}$. We always understand that $f(x) = 0$ if $x \notin \mathbb{N}$ for $f \in \mathcal{A}$. The set of all multiplicative and completely multiplicative arithmetic functions are respectively denoted by $\mathcal{M}$ and $\mathcal{M}^c$. Namely, $f \in \mathcal{M}$ (resp. $f \in \mathcal{M}^c$) means that $f(mn) = f(m)f(n)$ for $\gcd(m, n) = 1$ (resp. for all $m, n \in \mathbb{N}$). As usual, the product $fg \in \mathcal{A}$ of $f, g \in \mathcal{A}$ is defined by $(fg)(n) := f(n)g(n)$.

Let $\gamma \in \mathbb{N}$. We define the γ -convolution of $f, g \in \mathcal{A}$ by

$$(g *_{\gamma} f)(n) := \sum_{d^{\gamma} | n} f\left(\frac{n}{d^{\gamma}}\right) g(d^{\gamma}).$$

In particular, $*_1 = *$ stands for the usual Dirichlet convolution. Define the function $a_{\gamma} \in \mathcal{A}$ by $a_{\gamma}(n) = 1$ if there exists $d \in \mathbb{N}$ such that $n = d^{\gamma}$, and 0 otherwise. Then, it is clear that $g *_{\gamma} f = g^{[\gamma]} * f$, where $g^{[\gamma]} := a_{\gamma} g$. Note that the product $*_{\gamma}$ does not satisfy the commutativity or associativity properties unless $\gamma = 1$. We therefore inductively define the $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m)$ -convolution for $\gamma_1, \dots, \gamma_m \in \mathbb{N}$ of $f_1, \dots, f_{m+1} \in \mathcal{A}$ as

$$\begin{aligned} (f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1)(n) &:= ((f_{m+1} *_{\gamma_m} \cdots *_{\gamma_2} f_2) *_{\gamma_1} f_1)(n) \\ &= \sum_{d_m^{\gamma_m} | \cdots | d_1^{\gamma_1} | n} f_1\left(\frac{n}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) \cdots f_m\left(\frac{d_{m-1}^{\gamma_{m-1}}}{d_m^{\gamma_m}}\right) f_{m+1}(d_m^{\gamma_m}). \end{aligned}$$

Let $L(s; f) := \sum_{n=1}^{\infty} f(n) n^{-s}$ be the Dirichlet series attached to $f \in \mathcal{A}$ and let $\sigma(f) \in \mathbb{R} \cup \{\infty\}$ be the abscissa of absolute convergence of $L(s; f)$. Define $f^{(\gamma)} \in \mathcal{A}$ for $\gamma \in \mathbb{N}$ by $f^{(\gamma)}(n) := f(n^{\gamma})$. Then, it is clear that $L(\gamma s; f^{(\gamma)}) = L(s; f^{[\gamma]})$ with $\sigma(f^{[\gamma]}) \leq \sigma(f)$. For $f_1, \dots, f_{m+1} \in \mathcal{A}$, we here calculate the Dirichlet series attached to $f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1$.

Proposition 2.1. *Suppose $\gamma_0 | \gamma_1 | \gamma_2 | \cdots | \gamma_m$ with $\gamma_0 = 1$. Then, we have for $\operatorname{Re}(s) > \max\{\sigma(f_1), \dots, \sigma(f_{m+1})\}$*

$$L(s; f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1) = \prod_{j=1}^{m+1} L(s; f_j^{[\gamma_{j-1}]}). \quad (2.1)$$

Proof. Assume $d_m^{\gamma_m} | \cdots | d_1^{\gamma_1} | n$. Then, since $\gamma_1 | \gamma_2 | \cdots | \gamma_m$, we can write $d_j^{\gamma_j} = d_{j+1}^{\gamma_{j+1}} k_{j+1}^{\gamma_j}$ for $1 \leq j \leq m-1$ and $n = k_1 d_1^{\gamma_1} = \cdots = k_1 k_2^{\gamma_1} \cdots k_m^{\gamma_{m-1}} d_m^{\gamma_m}$ with $k_1, k_2, \dots, k_m \in \mathbb{N}$. We then write $d_m = k_{m+1}$, and so

$$\begin{aligned} L(s; f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1) &= \sum_{k_1, \dots, k_{m+1}=1}^{\infty} f_1(k_1) f_2(k_2^{\gamma_1}) \cdots f_{m+1}(k_{m+1}^{\gamma_m}) (k_1 k_2^{\gamma_1} \cdots k_{m+1}^{\gamma_m})^{-s} \\ &= L(s; f_1) L(\gamma_1 s; f_2^{(\gamma_1)}) \cdots L(\gamma_m s; f_{m+1}^{(\gamma_m)}). \end{aligned}$$

This ends the proof. $\square$

Remark 2.2. We cannot expect that $L(s; f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1)$ is expressed as a product of a Dirichlet series such as (2.1) for general $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m) \in \mathbb{N}^m$.

In the next subsection, we will introduce a multiple Ramanujan sum for $f_1, \dots, f_{m+1} \in \mathcal{A}$, which gives the $\boldsymbol{\gamma}$ -convolution $f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1$ in the diagonal case (for more detail, see Proposition 2.3 (iv)).

2.2 Definition of $S_{\mathbf{f}}^{\boldsymbol{\gamma}}$ and its basic properties

Let $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m) \in \mathbb{N}^m$ and $\mathbf{f} = (f_1, \dots, f_{m+1}) \in \mathcal{A}^{m+1}$. We define a multiple Ramanujan sum $S_{\mathbf{f}}^{\boldsymbol{\gamma}} = S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}$ of $\mathbf{f}$ with the parameter $\boldsymbol{\gamma}$ by

$$S_{\mathbf{f}}^{\boldsymbol{\gamma}}(n_1, \dots, n_{m+1}) := \sum_{\substack{d_j^{\gamma_j} | \gcd(n_1, \dots, n_{j+1}) \\ (j=1, \dots, m)}} f_1\left(\frac{n_1}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) \cdots f_m\left(\frac{d_{m-1}^{\gamma_{m-1}}}{d_m^{\gamma_m}}\right) f_{m+1}(d_m^{\gamma_m}),$$

where the sum is taken over all m -tuples $(d_1, \dots, d_m) \in \mathbb{N}^m$ satisfying $d_j^{\gamma_j} \mid \gcd(n_1, \dots, n_{j+1})$ for each $1 \leq j \leq m$. Note that the summand vanishes unless $d_m^{\gamma_m} \mid \dots \mid d_1^{\gamma_1} \mid n_1$. We write $S_{\mathbf{f}} := S_{\mathbf{f}}^{(1, \dots, 1)}$ and understand that $S_{\mathbf{f}}(n) = f(n)$ when $m = 0$. These are the fundamental properties of $S_{\mathbf{f}}^{\gamma}$, which are obtained from the elementary properties of the gcd-function.

Proposition 2.3. (i) For any $1 \leq j \leq m+1$, we have

$$S_{\mathbf{f}}^{\gamma}(n_1, \dots, n_{m+1}) = S_{f_1, \dots, f_{j-1}, S_{f_j, \dots, f_{m+1}}^{(\gamma_j, \dots, \gamma_m)}(\cdot, n_{j+1}, \dots, n_{m+1})}^{(\gamma_1, \dots, \gamma_{j-1})}(n_1, \dots, n_j). \quad (2.2)$$

(ii) For any $1 \leq j \leq m+1$, we have

$$S_{\mathbf{f}}^{\gamma}(n_1, \dots, n_{j-1}, 1, n_{j+1}, \dots, n_{m+1}) = f_j(1) \cdots f_{m+1}(1) \cdot S_{f_1, \dots, f_{j-1}}^{(\gamma_1, \dots, \gamma_{j-2})}(n_1, \dots, n_{j-1}).$$

In particular, we have $S_{\mathbf{f}}^{\gamma}(1, n_2, \dots, n_{m+1}) = f_1(1) \cdots f_{m+1}(1)$.

(iii) For any $1 \leq j \leq m$, we have

$$\begin{aligned} S_{f_1, \dots, f_{j-1}, \varepsilon, f_{j+1}, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}(n_1, \dots, n_{m+1}) \\ = S_{f_1, \dots, f_{j-1}, f_{j+1}, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_{j-1}, \gamma_{j+1}, \dots, \gamma_m)}(n_1, \dots, n_{j-1}, \gcd(n_j, n_{j+1}), n_{j+2}, \dots, n_{m+1}) \end{aligned} \quad (2.3)$$

and $S_{f_1, \dots, f_m, \varepsilon}^{(\gamma_1, \dots, \gamma_m)}(n_1, \dots, n_{m+1}) = S_{f_1, \dots, f_m}^{(\gamma_1, \dots, \gamma_{m-1})}(n_1, \dots, n_m)$.

(iv) Let $n_1 \mid n_j$ for all $2 \leq j \leq m+1$. Then, we have

$$S_{\mathbf{f}}^{\gamma}(n_1, \dots, n_{m+1}) = (f_{m+1} *_{\gamma_m} \cdots *_{\gamma_1} f_1)(n_1). \quad (2.4)$$

□

Example 2.4. Let $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{C}^m$. Define a multiple divisor function $\sigma_{\mathbf{a}}^{\gamma}$ with the parameter γ by

$$\sigma_{\mathbf{a}}^{\gamma}(n_1, \dots, n_{m+1}) := \sum_{\substack{d_m^{\gamma_m} \mid \dots \mid d_1^{\gamma_1} \mid n_1 \\ d_j^{\gamma_j} \mid \gcd(n_1, \dots, n_{j+1}) \ (j=1, \dots, m)}} d_1^{\gamma_1 a_1} \cdots d_m^{\gamma_m a_m}. \quad (2.5)$$

This is a generalization of the usual divisor function $\sigma_a(n) := \sum_{d \mid n} d^a$; $\sigma_a(n) = \sigma_a^{(1)}(n, n)$. Then, it holds that $\sigma_{\mathbf{a}}^{\gamma} = S_{\mathbf{f}}^{\gamma}$ with $f_j = \delta^{a_0 + a_1 + \dots + a_{j-1}}$ for $1 \leq j \leq m+1$ where $a_0 = 0$. Similarly, for $\mathbf{b} = (b_1, \dots, b_{m+1}) \in \mathbb{C}^{m+1}$, one can see that $\sigma_{\mathbf{b}}^{\gamma}(n_1, \dots, n_{m+1}) = n_1^{-b_1} S_{\delta^{b_1}, \dots, \delta^{b_{m+1}}}^{\gamma}(n_1, \dots, n_{m+1})$ where $\tilde{\mathbf{b}} := (b_2 - b_1, b_3 - b_2, \dots, b_{m+1} - b_m) \in \mathbb{C}^m$. Note that the sum $Z_n^{\gamma}(a_1, \dots, a_m) := \sigma_{\mathbf{a}}^{\gamma}(n, \dots, n)$ is studied in [11] and called the multiple finite Riemann zeta function. From formula (2.1), if $\gamma_0 \mid \gamma_1 \mid \gamma_2 \mid \dots \mid \gamma_m$ with $\gamma_0 = 1$, we have $L(s; Z^{\gamma}(a_1, \dots, a_m)) = \prod_{j=1}^{m+1} \zeta(\gamma_{j-1}(s - a_0 - \dots - a_{j-1}))$ where $\zeta(s) = L(s; \delta^0)$ is the Riemann zeta function.

Example 2.5. Let $f \in \mathcal{A}$. Then, the composition function $f \circ \gcd$ of f and the gcd-function can be expressed in terms of the multiple Ramanujan sum. Actually, from the degeneracy formula (2.3), we have $(f \circ \gcd)(n_1, \dots, n_{m+1}) = S_{\mathbf{f}}(n_1, \dots, n_{m+1})$ with $f_1 = \delta^0$, $f_2 = \dots = f_m = \varepsilon$ and $f_{m+1} = f * \mu$.

We next show the multiplicative property of the multiple Ramanujan sum S_f^γ . Recall that an arithmetic function $F(n_1, \dots, n_k)$ of k -variables is called multiplicative if

$$F(m_1 n_1, \dots, m_k n_k) = F(m_1, \dots, m_k) \cdot F(n_1, \dots, n_k)$$

for relatively prime k -tuples $(m_1, \dots, m_k) \in \mathbb{N}^k$ and $(n_1, \dots, n_k) \in \mathbb{N}^k$. Here, we say that $(m_1, \dots, m_k)$ and $(n_1, \dots, n_k)$ are relatively prime if $\gcd(m_i, n_j) = 1$ for all $1 \leq i, j \leq k$, and equivalently, $\gcd(n_1 \cdots n_k, m_1 \cdots m_k) = 1$ (see [21]). In this case, we have the following Euler product expression

$$F(n_1, \dots, n_k) = \prod_p F(p^{\alpha_{p,1}}, \dots, p^{\alpha_{p,k}}),$$

where $\alpha_{p,j} = \text{ord}_p n_j$ for $1 \leq j \leq k$. Note that this is a finite product since $F(1, \dots, 1) = 1$.

Proposition 2.6. *The function S_f^γ is multiplicative if $f_1, \dots, f_{m+1} \in \mathcal{M}$.*

Proof. Assume $(n_1, \dots, n_{m+1})$ and $(k_1, \dots, k_{m+1})$ are relatively prime. Then, for all $1 \leq j \leq m+1$, $\gcd(n_1 k_1, \dots, n_j k_j) = \gcd(n_1, \dots, n_j) \cdot \gcd(k_1, \dots, k_j)$ and $\gcd(n_1, \dots, n_j)$ and $\gcd(k_1, \dots, k_j)$ are relatively prime. Hence, the first assertion follows in the same manner as the that in the proof of Theorem 1 in [1]. $\square$

2.3 Finite Fourier expansions

An arithmetic function $F(r; n_1, \dots, n_k)$ of k -variables $n_1, \dots, n_k$ is called periodic (mod r) if $F(r; n_1, \dots, n_k) = F(r; n'_1, \dots, n'_k)$ whenever $n_j \equiv n'_j \pmod{r}$ for all $1 \leq j \leq k$ (see [13] for the case of $k = 1$). It is well-known that F is periodic (mod r) if and only if it has an expression of the form of

$$F(r; n_1, \dots, n_k) = \sum_{l_1, \dots, l_k=1}^r a_r(l_1, \dots, l_k) e(r, n_1 l_1) \cdots e(r, n_k l_k) \quad (2.6)$$

and the coefficients $a_r(l_1, \dots, l_k)$ are uniquely determined by

$$a_r(l_1, \dots, l_k) = \frac{1}{r^k} \sum_{n_1, \dots, n_k=1}^r F(r; n_1, \dots, n_k) e(r, -n_1 l_1) \cdots e(r, -n_k l_k). \quad (2.7)$$

Moreover, F is called even (mod r) if $F(r; n_1, \dots, n_k) = F(r; \gcd(n_1, r), \dots, \gcd(n_k, r))$. Note that F is periodic (mod r) if it is even (mod r). Then, as shown by E. Cohen [9], F is even (mod r) if and only if it has an expression of the form of

$$F(r; n_1, \dots, n_k) = \sum_{d_1, \dots, d_k | r} \alpha_r(d_1, \dots, d_k) c(d_1, n_1) \cdots c(d_k, n_k) \quad (2.8)$$

with

$$\alpha_r(d_1, \dots, d_k) = \frac{1}{r^k} \sum_{\delta_1, \dots, \delta_k | r} F(r; \delta_1, \dots, \delta_k) c\left(\frac{r}{\delta_1}, \frac{r}{d_1}\right) \cdots c\left(\frac{r}{\delta_k}, \frac{r}{d_k}\right). \quad (2.9)$$

We call expressions (2.6) and (2.8) finite Fourier expansions and coefficients a_r and α_r finite Fourier coefficients of F .

By definition, the multiple Ramanujan sum $S_{\mathbf{f}}^{\gamma}(n_1, \dots, n_{m+1})$ is even (mod n_1) as a function of m -variables $n_2, \dots, n_{m+1}$ for any $n_1 \in \mathbb{N}$. Then, let us calculate the finite Fourier expansions of $S_{\mathbf{f}}^{\gamma}$. To do so, we add one parameter (or, a weight) to $S_{\mathbf{f}}^{\gamma}$. Let ξ be an arithmetic function of m -variables. Set

$$\begin{aligned} S_{\mathbf{f}}^{\gamma, \xi}(n_1, \dots, n_{m+1}) &= S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m), \xi}(n_1, \dots, n_{m+1}) \\ &:= \sum_{\substack{d_j^{\gamma_j} \mid \gcd(n_1, \dots, n_{j+1}) \\ (j=1, \dots, m)}} \xi(d_1^{\gamma_1}, \dots, d_m^{\gamma_m}) f_1\left(\frac{n_1}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) \cdots f_m\left(\frac{d_{m-1}^{\gamma_{m-1}}}{d_m^{\gamma_m}}\right) f_{m+1}(d_m^{\gamma_m}). \end{aligned}$$

Here, $S_{\mathbf{f}}^{\gamma, \xi}(n_1, \dots, n_{m+1})$ is again even (mod n_1) as a function of $n_2, \dots, n_{m+1}$ for any $n_1 \in \mathbb{N}$ and $S_{\mathbf{f}}^{\gamma} = S_{\mathbf{f}}^{\gamma, 1_m}$ where $1_m(d_1, \dots, d_m) \equiv 1$. We also write $S_{\mathbf{f}}^{\xi} := S_{\mathbf{f}}^{(1, \dots, 1), \xi}$. As discussed in Proposition 2.6, $S_{\mathbf{f}}^{\gamma, \xi}$ can also be shown to be multiplicative if $f_1, \dots, f_{m+1}$ and ξ are multiplicative.

For $\mathbf{f} = (f_1, \dots, f_{m+1})$, we set ${}^t\tilde{\mathbf{f}} = (\delta^0 f_{m+1}, \delta^1 f_m, \dots, \delta^m f_1)$. Further, for $n \in \mathbb{N}$, we set

$${}^t\xi_n^{\gamma}(d_1, \dots, d_m) := \left(\prod_{j=1}^m a_{\gamma_j} \left(\frac{n}{d_{m+1-j}} \right) \right) \cdot \xi \left(\frac{n}{d_m}, \dots, \frac{n}{d_1} \right).$$

Then, we obtain the following theorem. Roughly speaking, the finite Fourier coefficients of $S_{\mathbf{f}}^{\gamma, \xi}$ can again be written as a multiple Ramanujan sum.

Theorem 2.7. *Write the finite Fourier expansions of $S_{\mathbf{f}}^{\gamma, \xi}$ as*

$$\begin{aligned} S_{\mathbf{f}}^{\gamma, \xi}(n_1, \dots, n_{m+1}) &= \sum_{l_2, \dots, l_{m+1}=1}^{n_1} a_{\mathbf{f}, n_1}^{\gamma, \xi}(l_2, \dots, l_{m+1}) e(n_1, n_2 l_2) \cdots e(n_1, n_{m+1} l_{m+1}) \\ &= \sum_{d_2, \dots, d_{m+1} \mid n_1} \alpha_{\mathbf{f}, n_1}^{\gamma, \xi}(d_2, \dots, d_{m+1}) c(d_2, n_2) \cdots e(d_{m+1}, n_{m+1}). \end{aligned}$$

Then, the finite Fourier coefficients $a_{\mathbf{f}, n_1}^{\gamma, \xi}$ and $\alpha_{\mathbf{f}, n_1}^{\gamma, \xi}$ of $S_{\mathbf{f}}^{\gamma, \xi}$ are respectively given as

$$a_{\mathbf{f}, n_1}^{\gamma, \xi}(l_2, \dots, l_{m+1}) = n_1^{-m} S_{t\tilde{\mathbf{f}}}^{t\xi_{n_1}^{\gamma}}(n_1, l_{m+1}, \dots, l_2), \quad (2.10)$$

$$\alpha_{\mathbf{f}, n_1}^{\gamma, \xi}(d_2, \dots, d_{m+1}) = n_1^{-m} S_{t\tilde{\mathbf{f}}}^{t\xi_{n_1}^{\gamma}}\left(n_1, \frac{n_1}{d_{m+1}}, \dots, \frac{n_1}{d_2}\right). \quad (2.11)$$

Proof. We only verify formula (2.11) (formula (2.10) can be similarly obtained). From formula (2.9), $\alpha_{\mathbf{f}, n_1}^{\gamma, \xi}(d_2, \dots, d_{m+1})$ is given as

$$n_1^{-m} \sum_{\delta_2, \dots, \delta_{m+1} \mid n_1} S_{\mathbf{f}}^{\gamma, \xi}(n_1, \delta_2, \dots, \delta_{m+1}) c\left(\frac{n_1}{\delta_2}, \frac{n_1}{d_2}\right) \cdots c\left(\frac{n_1}{\delta_{m+1}}, \frac{n_1}{d_{m+1}}\right).$$

Further, by changing the order of the summation, this expression is equivalent to

$$\begin{aligned} n_1^{-m} \sum_{e_m^{\gamma_m} \mid \cdots \mid e_1^{\gamma_1} \mid n_1} \xi(e_1^{\gamma_1}, \dots, e_m^{\gamma_m}) f_1\left(\frac{n_1}{e_1^{\gamma_1}}\right) f_2\left(\frac{e_1^{\gamma_1}}{e_2^{\gamma_2}}\right) \cdots f_m\left(\frac{e_{m-1}^{\gamma_{m-1}}}{e_m^{\gamma_m}}\right) f_{m+1}(e_m^{\gamma_m}) \\ \times \left(\sum_{e_1^{\gamma_1} \mid \delta_2 \mid n_1} c\left(\frac{n_1}{\delta_2}, \frac{n_1}{d_2}\right) \right) \cdots \left(\sum_{e_m^{\gamma_m} \mid \delta_{m+1} \mid n_1} c\left(\frac{n_1}{\delta_{m+1}}, \frac{n_1}{d_{m+1}}\right) \right). \quad (2.12) \end{aligned}$$

Here, we use the following identity. Let $e|n$ and $d|n$. Then, we have

$$\sum_{e|\delta|n} c\left(\frac{n}{\delta}, \frac{n}{d}\right) = \begin{cases} \frac{n}{e} & \text{if } d|e, \\ 0 & \text{otherwise.} \end{cases} \quad (2.13)$$

Actually, one can obtain this formula from the right-most expression (1.1) of the Ramanujan sum $c(k, n)$ and formula $\delta^0 * \mu = \varepsilon$. Then, applying formula (2.13), we see that (2.12) can be written as

$$\begin{aligned} n_1^{-m} \sum_{d_2|e_1^{\gamma_1}|n_1} \sum_{d_3|e_2^{\gamma_2}|e_1^{\gamma_1}} \cdots \sum_{d_{m+1}|e_m^{\gamma_m}|e_{m-1}^{\gamma_{m-1}}} \xi(e_1^{\gamma_1}, \dots, e_m^{\gamma_m}) \\ \times f_1\left(\frac{n_1}{e_1^{\gamma_1}}\right) f_2\left(\frac{e_1^{\gamma_1}}{e_2^{\gamma_2}}\right) \cdots f_m\left(\frac{e_{m-1}^{\gamma_{m-1}}}{e_m^{\gamma_m}}\right) f_{m+1}(e_m^{\gamma_m}) \frac{n_1}{e_1^{\gamma_1}} \cdots \frac{n_1}{e_m^{\gamma_m}}. \end{aligned}$$

Changing variables $e'_k = n_1/e_k^{\gamma_k}$ for all $1 \leq k \leq m$, we have $a_{\gamma_k}(n_1/e'_k) = 1$. Moreover, it holds that $e'_k | \frac{n_1}{d_{k+1}}$ for $1 \leq k \leq m$ and $e'_1 | \cdots | e'_m | n_1$, because $e_m^{\gamma_m} | \cdots | e_1^{\gamma_1} | n_1$. Hence, we can rewrite the above expression as

$$\begin{aligned} n_1^{-m} \sum_{\substack{e'_m | \frac{n_1}{d_{m+1}} \\ e'_m | \frac{n_1}{d_{m+1}}}} \sum_{\substack{e'_{m-1} | e'_m \\ e'_{m-1} | \frac{n_1}{d_m}}} \cdots \sum_{\substack{e'_1 | e'_2 \\ e'_1 | \frac{n_1}{d_2}}} a_{\gamma_1}\left(\frac{n_1}{e'_1}\right) \cdots a_{\gamma_m}\left(\frac{n_1}{e'_m}\right) \cdot \xi\left(\frac{n_1}{e'_1}, \dots, \frac{n_1}{e'_m}\right) \\ \times f_1(e'_1) f_2\left(\frac{e'_2}{e'_1}\right) \cdots f_m\left(\frac{e'_m}{e'_{m-1}}\right) f_{m+1}\left(\frac{n_1}{e'_m}\right) e'_1 \cdots e'_m. \end{aligned} \quad (2.14)$$

It is easy to see that (2.14) coincides with the right-hand side of formula (2.11). This completes the proof of theorem. $\square$

Remark 2.8. Suppose that $\gamma_0 | \gamma_1 | \cdots | \gamma_m$ with $\gamma_0 = 1$. Then, we have $S_{\tilde{\mathbf{f}}}^{t\xi_{n_1}^{\gamma}} = S_{t\tilde{\mathbf{f}}^{[\gamma]}}^{t\xi_{n_1}}$ where $t\xi_{n_1} := t\xi_{n_1}^{(1, \dots, 1)}$ and $t\tilde{\mathbf{f}}^{[\gamma]} = (\delta^0 f_{m+1}^{[\gamma_m]}, \delta^1 f_m^{[\gamma_{m-1}]}, \dots, \delta^m f_1^{[\gamma_0]})$. In fact, since the condition above means that $a_{\gamma_k}(n_1/e'_k) = a_{\gamma_k}(e'_{k+1}/e'_k)$ for all $1 \leq k \leq m-1$, the summand in (2.14) is written as

$$\xi\left(\frac{n_1}{e'_1}, \dots, \frac{n_1}{e'_m}\right) f_1(e'_1) f_2^{[\gamma_1]}\left(\frac{e'_2}{e'_1}\right) \cdots f_m^{[\gamma_{m-1}]}\left(\frac{e'_m}{e'_{m-1}}\right) f_{m+1}^{[\gamma_m]}\left(\frac{n_1}{e'_m}\right) e'_1 \cdots e'_m.$$

Hence, the claim follows.

Example 2.9. Retaining the notation in Example 2.4, one can easily see that $t\tilde{\mathbf{f}} = (\delta^{a_1+\cdots+a_m}, \delta^{a_1+\cdots+a_{m-1}+1}, \dots, \delta^{a_1+m-1}, \delta^m)$. Hence, the finite Fourier coefficient $\alpha_{\mathbf{f}, n_1} := \alpha_{\mathbf{f}, n_1}^{(1, \dots, 1), 1_m}$ of the multiple divisor function $\sigma_{\mathbf{a}} := \sigma_{\mathbf{a}}^{(1, \dots, 1)}$ is given by

$$\alpha_{\mathbf{f}, n_1}(d_2, \dots, d_{m+1}) = n_1^{a_1+\cdots+a_m-m} \sigma_{\mathbf{1}-t\mathbf{a}}\left(n_1, \frac{n_1}{d_{m+1}}, \dots, \frac{n_1}{d_2}\right),$$

where $\mathbf{1} := (1, \dots, 1) \in \mathbb{C}^m$ and $t\mathbf{a} := (a_m, \dots, a_1) \in \mathbb{C}^m$

Example 2.10. Retaining the notation in Example 2.5, we have $S_{\mathbf{f}} = f \circ \text{gcd}$. Again, by the degeneracy formula of (2.3), the finite Fourier coefficient $\alpha_{\mathbf{f}, n_1}$ of $f \circ \text{gcd}$ is given by

$$\alpha_{\mathbf{f}, n_1}(d_2, \dots, d_{m+1}) = n_1^{-m} S_{f * \mu, \delta^m}\left(n_1, \text{gcd}\left(\frac{n_1}{d_2}, \dots, \frac{n_1}{d_{m+1}}\right)\right).$$

3 Dirichlet series attached to S_f^γ

In this section, we examine both single-variable and multivariable Dirichlet series, the coefficients of which are given by the multiple Ramanujan sum S_f^γ .

3.1 Single-variable Dirichlet series

We first examine a single-variable Dirichlet series. Recall the following well-known formula concerning the Ramanujan sum $c(k, n)$ (see, e.g., [20]):

$$\sum_{k=1}^{\infty} c(k, n) k^{-s} = \frac{\sigma_{1-s}(n)}{\zeta(s)} \quad (\operatorname{Re}(s) > 1). \quad (3.1)$$

In this subsection, we give a generalization of this formula. For $j = 1, 2, \dots, m+1$, let

$$\Phi_f^\gamma(s; \check{n}_j) := \sum_{n_j=1}^{\infty} S_f^\gamma(n_1, \dots, n_{m+1}) n_j^{-s},$$

where $\check{n}_j := (n_1, \dots, n_{j-1}, n_{j+1}, \dots, n_{m+1}) \in \mathbb{N}^m$. The following proposition says that the series $\Phi_f^\gamma(s; \check{n}_j)$ is written as the product of a Dirichlet series and a finite sum, which is again given by the multiple Ramanujan sum.

Proposition 3.1. (i) For $j = 1$, we have

$$\Phi_f^\gamma(s; \check{n}_1) = L(s; f_1) S_{F_1}(n_2) = L(s; f_1) F_1(n_2) \quad (\operatorname{Re}(s) > \sigma(f_1)), \quad (3.2)$$

where $F_1 = F_{1, \check{n}_1}^{\gamma, s} := \delta^{-s} (S_{f_2, \dots, f_{m+1}}^{(\gamma_2, \dots, \gamma_m)}(\cdot, n_3, \dots, n_{m+1}) *_{\gamma_1} \delta^s)$.

(ii) For $2 \leq j \leq m+1$, we have

$$\Phi_f^\gamma(s; \check{n}_j) = \zeta(s) S_{f_1, \dots, f_{j-2}, F_j}^{(\gamma_1, \dots, \gamma_{j-2})}(n_1, \dots, n_{j-1}) \quad (\operatorname{Re}(s) > 1), \quad (3.3)$$

where $F_j = F_{j, \check{n}_j}^{\gamma, s} := \delta^{-s} (S_{f_j, \dots, f_{m+1}}^{(\gamma_j, \dots, \gamma_m)}(\cdot, n_{j+1}, \dots, n_{m+1}) *_{\gamma_{j-1}} (\delta^s f_{j-1}))$.

Proof. Suppose that $d_k^{\gamma k} \mid \gcd(n_1, \dots, n_{k+1})$ for all $k = 1, \dots, m$. Then, n_j is a multiple of $\operatorname{lcm}(d_1^{\gamma_1}, \dots, d_m^{\gamma_m}) = d_1^{\gamma_1}$ if $j = 1$ and $\operatorname{lcm}(d_{j-1}^{\gamma_{j-1}}, \dots, d_m^{\gamma_m}) = d_{j-1}^{\gamma_{j-1}}$ if $2 \leq j \leq m+1$ (note that we only consider m -tuples $(d_1, \dots, d_m)$ such that $d_m^{\gamma_m} \mid \dots \mid d_1^{\gamma_1}$). Hence, for $j = 1$ and $\operatorname{Re}(s) > \sigma(f_1)$, $\Phi_f^\gamma(s; \check{n}_j)$ is expressed as

$$\sum_{\substack{d_k^{\gamma k} \mid \gcd(n_2, \dots, n_{k+1}) \\ (k=1, \dots, m)}} \sum_{l=1}^{\infty} f_1\left(\frac{d_1^{\gamma_1} l}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) \cdots f_m\left(\frac{d_{m-1}^{\gamma_{m-1}}}{d_m^{\gamma_m}}\right) f_{m+1}(d_m^{\gamma_m}) (d_1^{\gamma_1} l)^{-s}$$

and, for $2 \leq j \leq m+1$ and $\operatorname{Re}(s) > 1$,

$$\begin{aligned} & \sum_{\substack{d_k^{\gamma k} \mid \gcd(n_1, \dots, n_{k+1}) \\ (k=1, 2, \dots, j-2)}} f_1\left(\frac{n_1}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) \cdots f_{j-2}\left(\frac{d_{j-3}^{\gamma_{j-3}}}{d_{j-2}^{\gamma_{j-2}}}\right) \sum_{d_{j-1}^{\gamma_{j-1}} \mid d_{j-2}^{\gamma_{j-2}}} f_{j-1}\left(\frac{d_{j-2}^{\gamma_{j-2}}}{d_{j-1}^{\gamma_{j-1}}}\right) \\ & \times \sum_{\substack{d_k^{\gamma k} \mid \gcd(d_{j-1}^{\gamma_{j-1}}, n_{j+1}, \dots, n_{k+1}) \\ (k=j, \dots, m)}} \sum_{l=1}^{\infty} f_j\left(\frac{d_{j-1}^{\gamma_{j-1}}}{d_j^{\gamma_j}}\right) \cdots f_m\left(\frac{d_{m-1}^{\gamma_{m-1}}}{d_m^{\gamma_m}}\right) f_{m+1}(d_m^{\gamma_m}) (d_{j-1}^{\gamma_{j-1}} l)^{-s}. \end{aligned}$$

Thus, it is easy to see that these coincide, respectively, with (3.2) and (3.3). This completes the proof. $\square$

Example 3.2. For small m , the series $\Phi_{\mathbf{f}}^{\gamma}(s; \check{\mathbf{n}}_j)$ is explicitly given as follows. For $m = 1$, we have

$$\Phi_{f_1, f_2}^{(\gamma_1)}(s; n_2) = L(s; f_1) \sum_{d^{\gamma_1} | n_2} f_2(d^{\gamma_1}) d^{-\gamma_1 s}, \quad (3.4)$$

$$\Phi_{f_1, f_2}^{(\gamma_1)}(s; n_1) = \zeta(s) \sum_{d^{\gamma_1} | n_1} f_1\left(\frac{n_1}{d^{\gamma_1}}\right) f_2(d^{\gamma_1}) d^{-\gamma_1 s} \quad (3.5)$$

and for $m = 2$, we have

$$\begin{aligned} \Phi_{f_1, f_2, f_3}^{(\gamma_1, \gamma_2)}(s; n_2, n_3) &= L(s; f_1) \sum_{d_1^{\gamma_1} | n_2} \sum_{d_2^{\gamma_2} | \gcd(d_1^{\gamma_1}, n_3)} f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) f_3(d_2^{\gamma_2}) d_1^{-\gamma_1 s}, \\ \Phi_{f_1, f_2, f_3}^{(\gamma_1, \gamma_2)}(s; n_1, n_3) &= \zeta(s) \sum_{d_1^{\gamma_1} | n_1} \sum_{d_2^{\gamma_2} | \gcd(d_1^{\gamma_1}, n_3)} f_1\left(\frac{n_1}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) f_3(d_2^{\gamma_2}) d_1^{-\gamma_1 s}, \\ \Phi_{f_1, f_2, f_3}^{(\gamma_1, \gamma_2)}(s; n_1, n_2) &= \zeta(s) \sum_{d_1^{\gamma_1} | \gcd(n_1, n_2)} \sum_{d_2^{\gamma_2} | d_1^{\gamma_1}} f_1\left(\frac{n_1}{d_1^{\gamma_1}}\right) f_2\left(\frac{d_1^{\gamma_1}}{d_2^{\gamma_2}}\right) f_3(d_2^{\gamma_2}) d_2^{-\gamma_2 s}. \end{aligned}$$

Formulas (3.4) and (3.5) were obtained in [1] for the case of $\gamma_1 = 1$.

Example 3.3. For $0 \leq k \leq m+1$ and $\mathbf{a} = (a_1, \dots, a_{m+1-k}) \in \mathbb{C}^{m+1-k}$, we define generalizations of the classical Ramanujan sum $c(n_1, n_2)$ as

$$c_{m+1, k}^{\mathbf{a}}(n_1, \dots, n_{m+1}) := S_{\underbrace{\mu, \dots, \mu}_{k}, \delta^{a_1}, \dots, \delta^{a_{m+1-k}}}(n_1, \dots, n_{m+1}).$$

It is clear that $c(n_1, n_2) = c_{2,1}^{(1)}(n_1, n_2)$. Then, for $1 \leq k \leq m$, using formulas (3.2) and $L(s; \mu) = 1/\zeta(s)$, we can verify by induction on m and k that

$$\begin{aligned} \sum_{n_1, \dots, n_k=1}^{\infty} c_{m+1, k}^{\mathbf{a}}(n_1, \dots, n_{m+1}) n_1^{-s_1} \dots n_k^{-s_k} \\ = \frac{\prod_{j=2}^k \zeta(s_j)}{\prod_{j=1}^k \zeta(s_1 + \dots + s_j)} \sum_{d | n_{k+1}} d^{a_1 - (s_1 + \dots + s_k)} \sigma_{\tilde{\mathbf{a}}}(d, n_{k+2}, \dots, n_{m+1}), \quad (3.6) \end{aligned}$$

where $\tilde{\mathbf{a}} := (a_2 - a_1, a_3 - a_2, \dots, a_{m+1-k} - a_{m-k}) \in \mathbb{C}^{m-k}$ (see Example 2.4). Setting $m = 1$, $k = 1$ (in this case, $\sigma_{\tilde{\mathbf{a}}} \equiv 1$) and $a_1 = 1$, we obtain formula (3.1). See [6] for a multivariable analogue of formula (3.1).

Example 3.4. Retaining the notation in Example 2.5, we have $S_{\mathbf{f}} = f \circ \gcd$. Let $\gcd(\check{\mathbf{n}}_j) := \gcd(n_1, \dots, n_{j-1}, n_{j+1}, \dots, n_{m+1})$. Then, since it holds that $\gcd(n_1, \dots, n_{m+1}) = \gcd(n_j, \gcd(\check{\mathbf{n}}_j))$, we have the following well-known formula

$$\sum_{n_j=1}^{\infty} (f \circ \gcd)(n_1, \dots, n_{m+1}) n_j^{-s} = \Phi_{\delta^0, f * \mu}(s; \gcd(\check{\mathbf{n}}_j)) = \zeta(s) \sum_{d | \gcd(\check{\mathbf{n}}_j)} (f * \mu)(d) d^{-s}.$$

3.2 Multivariable Dirichlet series

For an arithmetic function $F(n_1, \dots, n_k)$ of k -variables, we denote $L(\mathbf{s}; F)$ by the multi-variable Dirichlet series attached to F , that is,

$$L(\mathbf{s}; F) := \sum_{n_1, \dots, n_k=1}^{\infty} F(n_1, \dots, n_k) n_1^{-s_1} \cdots n_k^{-s_k}$$

for $\mathbf{s} = (s_1, \dots, s_k) \in \Sigma(F)$, where $\Sigma(F) \subset \mathbb{C}^k$ is the region of absolute convergence for $L(\mathbf{s}; F)$. The first goal of this subsection is to calculate $L(\mathbf{s}; S_{\mathbf{f}}^{\gamma})$ explicitly.

Theorem 3.5. *Let $\gamma_0 | \gamma_1 | \gamma_2 | \cdots | \gamma_m$ with $\gamma_0 \in \mathbb{N}$. Define the function $a_{\gamma_0} S_{\mathbf{f}}^{\gamma}$ of $(m+1)$ -variables as $(a_{\gamma_0} S_{\mathbf{f}}^{\gamma})(n_1, \dots, n_{m+1}) := a_{\gamma_0}(n_1) S_{\mathbf{f}}^{\gamma}(n_1, \dots, n_{m+1})$. Then, we have*

$$L(\mathbf{s}; a_{\gamma_0} S_{\mathbf{f}}^{\gamma}) = \prod_{j=2}^{m+1} \zeta(s_j) \cdot \prod_{j=1}^{m+1} L(s_1 + \cdots + s_j; f_j^{[\gamma_j-1]}) \quad (3.7)$$

in the region

$$\left\{ \mathbf{s} = (s_1, \dots, s_{m+1}) \in \mathbb{C}^{m+1} \left| \begin{array}{ll} \operatorname{Re}(s_j) > 1 & (2 \leq j \leq m+1), \\ \operatorname{Re}(s_1 + \cdots + s_j) > \sigma(f_j) & (1 \leq j \leq m+1) \end{array} \right. \right\}.$$

In particular, setting $\gamma_0 = 1$, we have

$$L(\mathbf{s}; S_{\mathbf{f}}^{\gamma}) = \prod_{j=2}^{m+1} \zeta(s_j) \cdot L(s_1; f_1) \prod_{j=2}^{m+1} L(s_1 + \cdots + s_j; f_j^{[\gamma_j-1]}). \quad (3.8)$$

Proof. This is proven by induction on m . Suppose that $m = 1$. Then, it holds from formula (3.5) that

$$\begin{aligned} L((s_1, s_2); a_{\gamma_0} S_{f_1, f_2}^{(\gamma_1)}) &= \sum_{n_1=1}^{\infty} \Phi_{f_1, f_2}^{(\gamma_1)}(s_2; n_1^{\gamma_0}) n_1^{-\gamma_0 s} \\ &= \zeta(s_2) \sum_{n_1=1}^{\infty} \sum_{d^{\gamma_1} | n_1^{\gamma_0}} f_1\left(\frac{n_1}{d^{\gamma_1}}\right) f_2(d^{\gamma_1}) d^{-\gamma_1 s_2} n_1^{-\gamma_0 s_1}. \end{aligned}$$

Here, since $\gamma_0 | \gamma_1$, $n_1^{\gamma_0}$ is expressed as $n_1^{\gamma_0} = d^{\gamma_1} l^{\gamma_0}$ with $l \in \mathbb{N}$. Hence, this expression is written as

$$\zeta(s_2) \sum_{d=1}^{\infty} \sum_{l=1}^{\infty} f_1\left(\frac{d^{\gamma_1} l^{\gamma_0}}{d^{\gamma_1}}\right) f_2(d^{\gamma_1}) d^{-\gamma_1 s_2} (d^{\gamma_1} l^{\gamma_0})^{-s_1},$$

thus completing the proof for $m = 1$. Next, suppose that $m-1$. Note that $S_{\mathbf{f}}^{\gamma}(n_1, \dots, n_{m+1}) = S_{f_1, S_{f_2, \dots, f_{m+1}}^{(\gamma_2, \dots, \gamma_m)}}^{(\gamma_1)}(\cdot, n_3, \dots, n_{m+1})(n_1, n_2)$ from formula (2.2). Then, from (3.7) for $m = 1$, $L(\mathbf{s}; a_{\gamma_0} S_{\mathbf{f}}^{\gamma})$ is equal to

$$\begin{aligned} & \sum_{n_3, \dots, n_{m+1}=1}^{\infty} L((s_1, s_2); a_{\gamma_0} S_{f_1, S_{f_2, \dots, f_{m+1}}^{(\gamma_2, \dots, \gamma_m)}}^{(\gamma_1)}(\cdot, n_3, \dots, n_{m+1})) n_3^{-s_1} \cdots n_{m+1}^{-s_{m+1}} \\ &= \zeta(s_2) L(s_1; f_1^{[\gamma_0]}) \sum_{n_3, \dots, n_{m+1}=1}^{\infty} L(s_1 + s_2; a_{\gamma_1} S_{f_2, \dots, f_{m+1}}^{(\gamma_2, \dots, \gamma_m)}(\cdot, n_3, \dots, n_{m+1})) n_3^{-s_1} \cdots n_{m+1}^{-s_{m+1}} \\ &= \zeta(s_2) L(s_1; f_1^{[\gamma_0]}) L((s_1 + s_2, s_3, \dots, s_{m+1}); a_{\gamma_1} S_{f_2, \dots, f_{m+1}}^{(\gamma_2, \dots, \gamma_m)}). \end{aligned}$$

This completes the proof for m based on the assumption of induction. Hence, we obtain the desired formula. $\square$

Remark that formula (2.1) is regarded as the “diagonally-summed version” of (3.8) since $S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)}(n, \dots, n) = (f_{m+1} *_{\gamma_m} \dots *_{\gamma_1} f_1)(n)$.

Example 3.6. Retaining the notation in Example 2.4, we have

$$L(\mathbf{s}; \sigma_{\mathbf{a}}) = \zeta(s_1) \prod_{k=2}^{m+1} \zeta(s_k) \zeta(s_1 + \dots + s_j - a_1 - \dots - a_{j-1}). \quad (3.9)$$

Example 3.7. Retaining the notation in Example 3.3, we have

$$L(\mathbf{s}; c_{m+1, k}^{\mathbf{a}}) = \frac{\prod_{l=2}^{m+1} \zeta(s_l) \prod_{l=1}^{m+1-k} \zeta(s_1 + \dots + s_{k+l} - a_l)}{\prod_{l=1}^k \zeta(s_1 + \dots + s_l)}.$$

We can also obtain this formula from (3.6) and (3.9). In particular, we have $L((s_1, s_2); c) = \zeta(s_2) \zeta(s_1 + s_2 - 1) / \zeta(s_1)$.

Example 3.8. Retaining the notation in Example 2.5, we have $S_{\mathbf{f}} = f \circ \gcd$. In this case, it holds that

$$L(\mathbf{s}; f \circ \gcd) = \frac{\zeta(s_1) \dots \zeta(s_{m+1})}{\zeta(s_1 + \dots + s_{m+1})} L(s_1 + \dots + s_{m+1}; f).$$

Next, we present another type of multivariable Dirichlet series involving $S_{\mathbf{f}}^{\gamma}$. Recall the well-known Ramanujan formula for the divisor function σ_a (see e.g., [20], also [7]):

$$L(s; \sigma_a \sigma_b) = \frac{\zeta(s) \zeta(s-a) \zeta(s-b) \zeta(s-a-b)}{\zeta(2s-a-b)}$$

for $\operatorname{Re}(s) > \max\{1, \operatorname{Re}(a) + 1, \operatorname{Re}(b) + 1, \operatorname{Re}(a+b) + 1\}$. This formula was generalized by J. M. Borwein and K. K. Choi [4] as follows. Let $f_1, f_2, g_1, g_2 \in \mathcal{M}^c$. Then, we have

$$L(s; (f_2 *_{\gamma} f_1)(g_2 *_{\gamma} g_1)) = \frac{L(s; f_1 g_1) L(s; (f_2 g_1)^{[\gamma]}) L(s; (f_1 g_2)^{[\gamma]}) L(s; (f_2 g_2)^{[\gamma]})}{L(2s; (f_1 f_2 g_1 g_2)^{[\gamma]})} \quad (3.10)$$

for $\operatorname{Re}(s) > \max\{\sigma(f_1 g_1), \sigma(f_2 g_1), \sigma(f_1 g_2), \sigma(f_2 g_2), \sigma(f_1 f_2 g_1 g_2)/2\}$. They gave formula (3.10) for the case in which $\gamma = 1$, and it is easy to obtain the equation for general $\gamma \in \mathbb{N}$ in the same manner. Similarly, since $S_{f_1, f_2}^{(\gamma)}(n, n) = (f_2 *_{\gamma} f_1)(n)$, we can regard formula (3.10) as the diagonally-summed version of the following formula.

Theorem 3.9. Let $f_1, f_2, g_1, g_2 \in \mathcal{M}^c$. Then, we have

$$\begin{aligned} & L((s_1, s_2); S_{f_1, f_2}^{(\gamma)} S_{g_1, g_2}^{(\gamma)}) \\ &= \frac{\zeta(s_2) L(s_1; f_1 g_1) L(s_1 + s_2; (f_2 g_1)^{[\gamma]}) L(s_1 + s_2; (f_1 g_2)^{[\gamma]}) L(s_1 + s_2; (f_2 g_2)^{[\gamma]})}{L(2(s_1 + s_2); (f_1 f_2 g_1 g_2)^{[\gamma]})} \end{aligned} \quad (3.11)$$

in the region

$$\left\{ (s_1, s_2) \in \mathbb{C}^2 \left| \begin{array}{l} \operatorname{Re}(s_2) > 1, \operatorname{Re}(s_1) > \sigma(f_1 g_1), \\ \operatorname{Re}(s_1 + s_2) > \max\{\sigma(f_2 g_1), \sigma(f_1 g_2), \sigma(f_2 g_2), \sigma(f_1 f_2 g_1 g_2)/2\} \end{array} \right. \right\}.$$

Proof. First, note that, for a multiplicative arithmetic function F of k -variables, $L(\mathbf{s}; F)$ has the Euler product expression $L(\mathbf{s}; F) = \prod_p \sum_{l_1, \dots, l_k=1}^{\infty} F(p^{l_1}, \dots, p^{l_k}) p^{-s_1 l_1} \dots p^{-s_k l_k}$.

Let $h_1, h_2 \in \mathcal{M}^c$. Then, it holds that

$$S_{h_1, h_2}^{(\gamma)}(p^{l_1}, p^{l_2}) = \frac{h_1(p)^\gamma}{h_1(p)^\gamma - h_2(p)^\gamma} h_1(p)^{l_1} \left(1 - \left(\frac{h_2(p)}{h_1(p)} \right)^{\gamma \cdot m^\gamma(l_1, l_2)} \right),$$

where $m^\gamma(l_1, l_2) := \lfloor \frac{1}{\gamma} \min\{l_1, l_2\} \rfloor + 1$. Hence, from the Euler product expression, we have

$$\begin{aligned} & L((s_1, s_2); S_{f_1, f_2}^{(\gamma)} S_{g_1, g_2}^{(\gamma)}) \\ &= \prod_p \sum_{l_1, l_2=0}^{\infty} S_{f_1, f_2}^{(\gamma)}(p^{l_1}, p^{l_2}) S_{g_1, g_2}^{(\gamma)}(p^{l_1}, p^{l_2}) p^{-s_1 l_1} p^{-s_2 l_2} \\ &= \prod_p (C_p^\gamma)^{-1} \sum_{l_1, l_2=0}^{\infty} (a_1 b_1 x_1)^{l_1} x_2^{l_2} \left(1 - \left(\frac{a_2}{a_1} \right)^{\gamma \cdot m^\gamma(l_1, l_2)} \right) \left(1 - \left(\frac{b_2}{b_1} \right)^{\gamma \cdot m^\gamma(l_1, l_2)} \right), \end{aligned} \quad (3.12)$$

where $a_i := f_i(p)$, $b_i := g_i(p)$, $x_i := p^{-s_i}$ for $i = 1, 2$ and $C_p^\gamma = C_p^\gamma(f_1, f_2, g_1, g_2) := (a_1^\gamma - a_2^\gamma)(b_1^\gamma - b_2^\gamma)/(a_1 b_1)^\gamma$ for each prime p . Write I for the inner sum of the right-most hand side of (3.12). Moreover, divide I into two parts as $\sum_{l_1, l_2=0}^{\infty} = \sum_{l_1 \geq l_2}^{\infty} + \sum_{l_1 < l_2}^{\infty} = \sum_{l_2=0}^{\infty} \sum_{l_1=l_2}^{\infty} + \sum_{l_1=0}^{\infty} \sum_{l_2=l_1+1}^{\infty}$ and denote I_1 and I_2 by the former and latter sums, respectively. Writing l_2 as $l_2 = \gamma l + k$ in I_1 and l_1 as $l_1 = \gamma l + k$ in I_2 with $l \in \mathbb{Z}_{\geq 0}$ and $0 \leq k \leq \gamma - 1$, respectively, we have

$$\begin{aligned} I_1 &= \sum_{k=0}^{\gamma-1} \sum_{l=0}^{\infty} \sum_{l_1=\gamma l+k}^{\infty} (a_1 b_1 x_1)^{l_1} x_2^{\gamma l+k} \left(1 - \left(\frac{a_2}{a_1} \right)^{\gamma(l+1)} \right) \left(1 - \left(\frac{b_2}{b_1} \right)^{\gamma(l+1)} \right) \\ &= \frac{1}{1 - a_1 b_1 x_1} \frac{1 - (a_1 b_1 x_1 x_2)^\gamma}{1 - a_1 b_1 x_1 x_2} M \end{aligned}$$

and similarly

$$I_2 = \frac{x_2}{1 - x_2} \frac{1 - (a_1 b_1 x_1 x_2)^\gamma}{1 - a_1 b_1 x_1 x_2} M,$$

where $M := \sum_{l=0}^{\infty} (a_1 b_1 x_1 x_2)^{\gamma l} \left(1 - \left(\frac{a_2}{a_1} \right)^{\gamma(l+1)} \right) \left(1 - \left(\frac{b_2}{b_1} \right)^{\gamma(l+1)} \right)$. Hence, we have

$$I = I_1 + I_2 = \frac{1 - (a_1 b_1 x_1 x_2)^\gamma}{(1 - x_2)(1 - a_1 b_1 x_1)} M.$$

Now, M is straightforwardly calculated as

$$\frac{C_p^\gamma (1 - (a_1 a_2 b_1 b_2 x_1^2 x_2^2)^\gamma)}{(1 - (a_1 b_1 x_1 x_2)^\gamma) (1 - (a_2 b_1 x_1 x_2)^\gamma) (1 - (a_1 b_2 x_1 x_2)^\gamma) (1 - (a_2 b_2 x_1 x_2)^\gamma)},$$

where the desired formula immediately follows from formula (3.12). $\square$

Remark 3.10. For $m \geq 2$, we cannot expect that the single-variable Dirichlet series $L(s; (f_{m+1} *_{\gamma_m} \dots *_{\gamma_1} f_1)(g_{m+1} *_{\gamma_m} \dots *_{\gamma_1} g_1))$ has a product expression such as (3.10) and, similarly, the multivariable Dirichlet series $L((s_1, \dots, s_{m+1}); S_{f_1, \dots, f_{m+1}}^{(\gamma_1, \dots, \gamma_m)} S_{g_1, \dots, g_{m+1}}^{(\gamma_1, \dots, \gamma_m)})$ has a product expression such as (3.11).

4 A generalization of the Smith determinant

In this section, we evaluate the hyperdeterminants in the sense of Cayley of the hypermatrices, the entries of which are given by values of the multiple Ramanujan sum $S_f^{\gamma, \xi}$.

4.1 Hyperdeterminant

Recall the definition and some properties of the hyperdeterminants in the sense of Cayley. Let $A = (A(i_1, \dots, i_k))_{1 \leq i_1, \dots, i_k \leq n}$ be a k -dimensional matrix of order n . For a subset $I \subseteq \{1, \dots, k\}$, set $\varepsilon_j = 1$ if $j \in I$, and 0 otherwise. Then, the hyperdeterminant $\det_I A$ of A with the signature I is defined as

$$\det_I A := \frac{1}{n!} \sum_{\sigma_1, \dots, \sigma_k \in \mathfrak{S}_n} \prod_{j=1}^k \text{sgn}(\sigma_j)^{\varepsilon_j} \sum_{v=1}^n A(\sigma_1(v), \dots, \sigma_k(v)), \quad (4.1)$$

where $\mathfrak{S}_n$ is the symmetric group of degree n . We obtain the usual determinant of a matrix A when $k = 2$ and $I = \{1, 2\}$; $\det_{\{1, 2\}} A = \det A$. Note that $\det_I A$ is identically zero if the number of elements in I is odd. For the theory of hyperdeterminants, see, e.g., [16]. We next present some lemmas that will be needed in the subsequent discussion.

Lemma 4.1. *Let $A = (A(i_1, \dots, i_k))$ be a k -dimensional matrix of order n .*

(i) *For $\pi \in \mathfrak{S}_n$, we have*

$$\det_I (A(\pi(i_1), \dots, \pi(i_k))) = \det_I (A(i_1, \dots, i_k)). \quad (4.2)$$

(ii) *For $\pi \in \mathfrak{S}_k$, set $\pi^{-1}(I) := \{\pi^{-1}(i) \mid i \in I\}$. Then, we have*

$$\det_I (A(i_{\pi(1)}, \dots, i_{\pi(k)})) = \det_{\pi^{-1}(I)} (A(i_1, \dots, i_k)). \quad (4.3)$$

Proof. These are immediately obtained from the definition of the hyperdeterminant. $\square$

Let $A = (A(i_1, \dots, i_k))$ and $B = (B(i_1, \dots, i_l))$ be k -dimensional and l -dimensional matrices, respectively, of order n . Then, the Cayley product AB of A and B is a $(k+l-2)$ -dimensional matrix of order n given by

$$(AB)(i_1, \dots, i_{k+l-2}) := \sum_{j=1}^n A(i_1, \dots, i_{k-1}, j) B(j, i_k, \dots, i_{k+l-2}). \quad (4.4)$$

Lemma 4.2. *Let $A = (A(i_1, \dots, i_k))$ and $B = (B(i_1, \dots, i_l))$ be k -dimensional and l -dimensional matrices, respectively, of order n . Let $K \subseteq \{1, 2, \dots, k-1\}$ and $L \subseteq \{2, 3, \dots, l\}$ with odd cardinality. Set $I := K \cup (L + (k-2))$, where $L + (k-2) := \{l+k-2 \mid l \in L\}$. Then, we have*

$$\det_I (AB) = \det_{K \cup \{k\}} A \cdot \det_{\{1\} \cup L} B. \quad (4.5)$$

Proof. See [10, 18]. $\square$

4.2 Smith hyperdeterminants

Let $S = \{x_1, \dots, x_n\}$ be a set of distinct positive integers. The goal of this subsection is to evaluate the hyperdeterminant $\det_I(S_{\mathbf{f}}^{\gamma, \xi}(x_{i_1}, \dots, x_{i_{m+1}}))$ for a factor-closed set S . Here, S is called factor-closed if S contains every divisor of x for any $x \in S$. To accomplish this, we slightly extend the definition of the even arithmetic function examined in Section 2.

First, we define some notations. For $\mathbf{d} = (d_1, \dots, d_k) \in \mathbb{N}^k$, we set $|\mathbf{d}| := \sum_{j=1}^k d_j$ and $x_{\mathbf{d}} := (x_{d_1}, \dots, x_{d_k})$. Moreover, for $r \in \mathbb{N}$, we set $\gcd(\mathbf{d}, r) := (\gcd(d_1, r), \dots, \gcd(d_k, r)) \in \mathbb{N}^k$, write $\mathbf{d}|r$ if $d_j|r$ for $1 \leq j \leq k$ and, in this case, set $r/\mathbf{d} := (r/d_1, \dots, r/d_k) \in \mathbb{N}^k$. Let $f(m, n)$ be an arithmetic function of 2-variables. For $\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}^k$ and $\mathbf{n} = (n_1, \dots, n_k) \in \mathbb{N}^k$, we set $f(\mathbf{n}, \mathbf{m}) := \prod_{j=1}^k f(m_j, n_j)$.

Let $\mathbf{r} = (r_1, \dots, r_m) \in \mathbb{N}^m$ and $\mathbf{k} = (k_1, \dots, k_m) \in \mathbb{N}^m$. Let $\mathbf{n}_j = (n_{j,1}, \dots, n_{j,k_j}) \in \mathbb{N}^{k_j}$ for $j = 1, \dots, m$. We call an arithmetic function $F(\mathbf{r}; \mathbf{n}_1, \dots, \mathbf{n}_m)$ of $|\mathbf{k}|$ -variables $\mathbf{n}_1, \dots, \mathbf{n}_m$ even (mod $\mathbf{r}^{(\mathbf{k})}$) if it holds that

$$F(\mathbf{r}; \gcd(\mathbf{n}_1, r_1), \dots, \gcd(\mathbf{n}_m, r_m)) = F(\mathbf{r}; \mathbf{n}_1, \dots, \mathbf{n}_m).$$

Then, the following lemma is obtained as described in [9] (note that the case of $m = 1$ is nothing more than Theorem 22 in [9]).

Lemma 4.3. *A function $F(\mathbf{r}; \mathbf{n}_1, \dots, \mathbf{n}_m)$ is even (mod $\mathbf{r}^{(\mathbf{k})}$) if and only if it possesses a representation of the form*

$$F(\mathbf{r}; \mathbf{n}_1, \dots, \mathbf{n}_m) = \sum_{\substack{\mathbf{d}_j | r_j \\ (j=1, \dots, m)}} \alpha_{\mathbf{r}}(\mathbf{d}_1, \dots, \mathbf{d}_m) \prod_{j=1}^m c(\mathbf{d}_j, \mathbf{n}_j), \quad (4.6)$$

where the sum is taken over all $\mathbf{d}_j = (d_{j,1}, \dots, d_{j,k_j}) \in \mathbb{N}^{k_j}$ for $j = 1, \dots, m$ satisfying $\mathbf{d}_j | r_j$. Moreover, the finite Fourier coefficients $\alpha_{\mathbf{r}}(\mathbf{d}_1, \dots, \mathbf{d}_m)$ are uniquely determined by

$$\alpha_{\mathbf{r}}(\mathbf{d}_1, \dots, \mathbf{d}_m) = \frac{1}{r_1^{k_1} \dots r_m^{k_m}} \sum_{\substack{\boldsymbol{\delta}_j | r_j \\ (j=1, \dots, m)}} F(\mathbf{r}; \boldsymbol{\delta}_1, \dots, \boldsymbol{\delta}_m) \prod_{j=1}^m c\left(\frac{r_j}{\boldsymbol{\delta}_j}, \frac{r_j}{\mathbf{d}_j}\right), \quad (4.7)$$

where the sum is also taken over all $\boldsymbol{\delta}_j = (\delta_{j,1}, \dots, \delta_{j,k_j}) \in \mathbb{N}^{k_j}$ for $j = 1, \dots, m$ satisfying $\boldsymbol{\delta}_j | r_j$. $\square$

Now, we obtain the following proposition.

Proposition 4.4. *Let $S = \{x_1, \dots, x_n\}$ be a factor-closed set, and let $F(\mathbf{r}; \mathbf{n}_1, \dots, \mathbf{n}_k)$ be an even function (mod $\mathbf{r}^{(\mathbf{k})}$). Define the $(m + |\mathbf{k}|)$ -dimensional matrix B of order n as*

$$B(i_1, \dots, i_{m+|\mathbf{k}|}) = F(x_{i_1}, \dots, x_{i_m}; \underbrace{x_{i_{m+1}}, \dots, x_{i_{m+k_1}}}_{k_1}, \dots, \underbrace{x_{i_{m+k_1+\dots+k_{m-1}+1}}, \dots, x_{i_{m+|\mathbf{k}|}}}_{k_m}).$$

Let I be a subset of $\{1, 2, \dots, m + |\mathbf{k}|\}$ with even cardinality such that $\{m+1, m+2, \dots, m+|\mathbf{k}|\} \subseteq I$, and set $\varepsilon_j = 1$ if $j \in I$, and 0 otherwise. Define the subset $\tilde{I}$ of $\{1, 2, \dots, m\}$ as $j \in \tilde{I}$ if and only if $\varepsilon_j + k_j$ is odd for $1 \leq j \leq m$. Then, we have

$$\det_I B = (x_1 \dots x_n)^{|\mathbf{k}|} \det_{\tilde{I}}(\alpha_{x_{i_1}, \dots, x_{i_m}}(\underbrace{x_{i_1}, \dots, x_{i_{k_1}}}_{k_1}, \dots, \underbrace{x_{i_m}, \dots, x_{i_{k_m}}}_{k_m}))_{1 \leq i_1, \dots, i_m \leq n}. \quad (4.8)$$

Here, $\alpha_{x_{i_1}, \dots, x_{i_m}}$ is the finite Fourier coefficient of F given by (4.7).

We then need the following lemma.

Lemma 4.5. *Let $A = (A(i_1, \dots, i_k))$ and $C = (C(i_1, i_2))$ be k -dimensional and 2-dimensional matrices, respectively, of order n . For $l = 0, 1, \dots, k$, define the k -dimensional matrices $A_C^{(l)}$ of order n by the following recursion formula:*

$$\begin{cases} A_C^{(0)}(i_1, \dots, i_k) := A(i_1, \dots, i_k) & (l = 0), \\ A_C^{(l)}(i_1, \dots, i_k) := (A_C^{(l-1)}C)(i_2, \dots, i_k, i_1) & (l = 1, 2, \dots, k), \end{cases}$$

where $A_C^{(l-1)}C$ is the Cayley product of $A_C^{(l-1)}$ and C . Then, we have

$$A_C^{(l)}(i_1, \dots, i_k) = \sum_{j_1, \dots, j_l=1}^n A(i_{l+1}, \dots, i_k, j_1, \dots, j_l) \prod_{h=1}^l C(j_h, i_h). \quad (4.9)$$

Proof. This is shown by induction on l from the definition of the Cayley product (4.4). $\square$

Proof of Proposition 4.4. From formula (4.2), we can assume that $x_1 < x_2 < \dots < x_n$. Define the $(m + |\mathbf{k}|)$ -dimensional matrix A and 2-dimensional matrices C of order n , respectively, as

$$A(i_1, \dots, i_{m+|\mathbf{k}|}) := \alpha_{x_{i_1}, \dots, x_{i_m}}(x_{i_{m+1}}, \dots, x_{i_{m+|\mathbf{k}|}}) \prod_{j=1}^m \prod_{l_j=1}^{k_j} e_{i_{m+k_1+\dots+k_{j-1}+l_j}, i_j},$$

$$C(i_1, i_2) := c(x_{i_1}, x_{i_2}),$$

where $e_{j,i} = 1$ if $x_j | x_i$, and 0 otherwise. Let $\rho = (1, 2, \dots, m + |\mathbf{k}|) \in \mathfrak{S}_{m+|\mathbf{k}|}$ be the cyclic permutation of order $(m + |\mathbf{k}|)$. Then, we have

$$B(i_1, \dots, i_{m+|\mathbf{k}|}) = A_C^{(|\mathbf{k}|)}(i_{\rho^m(1)}, \dots, i_{\rho^m(m+|\mathbf{k}|)}). \quad (4.10)$$

In fact, by equation (4.9), the right-hand side of (4.10) is equal to

$$\begin{aligned} & A_C^{(|\mathbf{k}|)}(i_{m+1}, \dots, i_{m+|\mathbf{k}|}, i_1, \dots, i_m) \\ &= \sum_{\substack{d_{j,1}, \dots, d_{j,k_j}=1 \\ (j=1, \dots, m)}}^n \alpha_{x_{i_1}, \dots, x_{i_m}} \left(\underbrace{x_{d_{1,1}}, \dots, x_{d_{1,k_1}}}_{k_1}, \dots, \underbrace{x_{d_{m,1}}, \dots, x_{d_{m,k_m}}}_{k_m} \right) \\ & \quad \times \prod_{j=1}^m \prod_{l_j=1}^{k_j} e_{d_{j,l_j}, i_j} c(x_{d_{j,l_j}}, x_{i_{m+k_1+\dots+k_{j-1}+l_j}}) \\ &= \sum_{\substack{\mathbf{d}_j | x_{i_j} \\ (j=1, \dots, m)}} \alpha_{x_{i_1}, \dots, x_{i_m}}(x_{\mathbf{d}_1}, \dots, x_{\mathbf{d}_m}) \prod_{j=1}^m \prod_{l_j=1}^{k_j} c(x_{d_{j,l_j}}, x_{i_{m+k_1+\dots+k_{j-1}+l_j}}), \end{aligned}$$

where $\mathbf{d}_j := (d_{j,1}, \dots, d_{j,k_j})$ for $1 \leq j \leq m$. This clearly coincides with the left-hand side of formula (4.10) from the expression of (4.6) of F , thus completing the proof. Note that,

since S is factor-closed, every divisor of x_j can again be written as x_d for some d . Next, using formula (4.3), we have

$$\begin{aligned}\det_I B &= \det_I (A_C^{(|\mathbf{k}|)}(i_{\rho^m(1)}, \dots, i_{\rho^m(m+|\mathbf{k}|)})) \\ &= \det_I ((A_C^{(|\mathbf{k}|-1)}C)(i_{\rho^m(2)}, \dots, i_{\rho^m(m+|\mathbf{k}|)}, i_{\rho^m(1)})) \\ &= \det_I ((A_C^{(|\mathbf{k}|-1)}C)(i_{\rho^{m+1}(1)}, \dots, i_{\rho^{m+1}(m+|\mathbf{k}|)})) \\ &= \det_{\rho^{-m-1}(I)} ((A_C^{(|\mathbf{k}|-1)}C)(i_1, \dots, i_{m+|\mathbf{k}|})).\end{aligned}$$

Here, note that $m+1 \in I$ implies $m+|\mathbf{k}| = \rho^{-m-1}(m+1) \in \rho^{-m-1}(I)$. Therefore, using formula (4.5) with $K = \rho^{-m-1}(I) \setminus \{m+|\mathbf{k}|\}$ and $L = \{2\}$, we see that the right-hand side of the above equation is equivalent to

$$\begin{aligned}\det_{(\rho^{-m-1}(I) \setminus \{m+|\mathbf{k}|\}) \cup \{m+|\mathbf{k}|\}} (A_C^{(|\mathbf{k}|-1)}) \cdot \det_{\{1,2\}} C \\ = \det C \cdot \det_{\rho^{-m-1}(I)} ((A_C^{(|\mathbf{k}|-2)}C)(i_2, \dots, i_{m+|\mathbf{k}|}, i_1)) \\ = \det C \cdot \det_{\rho^{-m-2}(I)} ((A_C^{(|\mathbf{k}|-2)}C)(i_1, \dots, i_{m+|\mathbf{k}|})).\end{aligned}$$

Under the condition $m+1, \dots, m+|\mathbf{k}| \in I$, the above procedure yields

$$\det_I B = (\det_{\{1,2\}} C)^{|\mathbf{k}|} \cdot \det_{\rho^{-m-|\mathbf{k}|}(I)} (A_C^{(0)}) = (\det_{\{1,2\}} C)^{|\mathbf{k}|} \cdot \det_I A.$$

By Corollary 3 in [5], we have $\det C = \det(c(x_{i_1}, x_{i_2})) = x_1 \cdots x_n$. Hence, it is sufficient to calculate $\det_I A$. Since $\varepsilon_j = 1$ for $j = m+1, \dots, m+|\mathbf{k}|$, we have

$$\begin{aligned}\det_I A &= \frac{1}{n!} \sum_{\boldsymbol{\sigma}} \sum_{\sigma_1, \dots, \sigma_m} \prod_{j=1}^m \text{sgn}(\sigma_j)^{\varepsilon_j} \prod_{j=1}^m \prod_{l_j=1}^{k_j} \text{sgn}(\sigma_{j,l_j}) \\ &\quad \times \prod_{v=1}^n \alpha_{x_{\boldsymbol{\sigma}(v)}}(x_{\sigma_1(v)}, \dots, x_{\sigma_m(v)}) \prod_{j=1}^m \prod_{l_j=1}^{k_j} e_{\sigma_{j,l_j}(v), \sigma_j(v)}.\end{aligned}$$

Here, the sum is taken over all $\boldsymbol{\sigma} = (\sigma_1, \dots, \sigma_m) \in (\mathfrak{S}_n)^m$ and $\sigma_j = (\sigma_{j,1}, \dots, \sigma_{j,k_j}) \in (\mathfrak{S}_n)^{k_j}$ for $1 \leq j \leq m$. For $\boldsymbol{\sigma} = (\sigma_1, \dots, \sigma_k) \in (\mathfrak{S}_n)^k$ and $v \in \{1, \dots, n\}$, we set $\boldsymbol{\sigma}(v) := (\sigma_1(v), \dots, \sigma_k(v))$. Replacing the variables $\tau_{j,l_j} = \sigma_{j,l_j} \sigma_j^{-1}$ with $1 \leq j \leq m$ and $1 \leq l_j \leq k_j$ and writing $\boldsymbol{\tau}_j = (\tau_{j,1}, \dots, \tau_{j,k_j})$, we have

$$\begin{aligned}\det_I A &= \frac{1}{n!} \sum_{\boldsymbol{\sigma}} \prod_{j=1}^m \text{sgn}(\sigma_j)^{\varepsilon_j + k_j} \sum_{\boldsymbol{\tau}_1, \dots, \boldsymbol{\tau}_m} \prod_{j=1}^m \prod_{l_j=1}^{k_j} \text{sgn}(\tau_{j,l_j}) \\ &\quad \times \prod_{v=1}^n \alpha_{x_{\boldsymbol{\sigma}(v)}}(x_{\boldsymbol{\tau}_1 \sigma_1(v)}, \dots, x_{\boldsymbol{\tau}_m \sigma_m(v)}) \prod_{j=1}^m \prod_{l_j=1}^{k_j} e_{\tau_{j,l_j} \sigma_j(v), \sigma_j(v)},\end{aligned}$$

where $\boldsymbol{\tau}_j \sigma_j = (\tau_{j,1} \sigma_j, \dots, \tau_{j,k_j} \sigma_j)$ for $1 \leq j \leq m$. Note that $e_{\tau_{j,l_j} \sigma_j(v), \sigma_j(v)} = 1$ holds for all $1 \leq v \leq n$ if and only if $\tau_{j,l_j} = 1$ since $x_1 < x_2 < \dots < x_n$. Therefore, $\det_I A$ is equal to

$$\begin{aligned}\frac{1}{n!} \sum_{\boldsymbol{\sigma}} \prod_{j=1}^m \text{sgn}(\sigma_j)^{\varepsilon_j + k_j} \prod_{v=1}^n \alpha_{x_{\boldsymbol{\sigma}(v)}}(x_{\sigma_1(v)}, \dots, x_{\sigma_m(v)}) \\ = \det_{\tilde{I}} (\alpha_{x_{i_1}, \dots, x_{i_m}} (\underbrace{x_{i_1}, \dots, x_{i_1}}_{k_1}, \dots, \underbrace{x_{i_m}, \dots, x_{i_m}}_{k_m}))_{1 \leq i_1, \dots, i_m \leq n}.\end{aligned}$$

This completes the proof of the proposition. $\square$

Theorem 4.6. Let $S = \{x_1, \dots, x_n\}$ be a factor-closed set and let $F(r; n_1, \dots, n_k)$ be an even function (mod r) with respect to the variables $n_1, \dots, n_k$. Let $I = \{2, 3, \dots, k+1\}$ if k is even, and $\{1, 2, \dots, k+1\}$ otherwise. Then, we have

$$\det_I(F(x_{i_1}; x_{i_2}, \dots, x_{i_{k+1}}))_{1 \leq i_1, \dots, i_{k+1} \leq n} = (x_1 \cdots x_n)^k \prod_{v=1}^n \alpha_{x_v}(x_v, \dots, x_v), \quad (4.11)$$

where α_{x_v} is the finite Fourier coefficient given by (2.9).

Proof. Setting $m = 1$ in Proposition 4.4, we immediately obtain this theorem. $\square$

Remark 4.7. The case of $k = 1$ is obtained in Theorem 2 in [5].

Corollary 4.8. Let $S = \{x_1, \dots, x_n\}$ be a factor-closed set. Let $I = \{2, 3, \dots, m+1\}$ if m is even, and $\{1, 2, \dots, m+1\}$ otherwise. Let ξ be arbitrary arithmetic function of m -variables. Then, we have

$$\begin{aligned} \det_I(S_{\mathbf{f}}^{\gamma, \xi}(x_{i_1}, \dots, x_{i_{m+1}}))_{1 \leq i_1, \dots, i_{m+1} \leq n} \\ = (f_1(1) \cdots f_m(1))^n \prod_{v=1}^n \xi(x_v, \dots, x_v) f_{m+1}^{[\text{lcm}(\gamma_1, \dots, \gamma_m)]}(x_v). \end{aligned} \quad (4.12)$$

Proof. From formula (2.11), we have $\alpha_{\mathbf{f}, x_v}^{\gamma, \xi}(x_v, \dots, x_v) = x_v^{-m} S_{t, \mathbf{f}}^{t, \xi, \gamma_v}(x_v, 1, \dots, 1)$. Furthermore, using the identity $a_{\gamma_1}(x_v) \cdots a_{\gamma_m}(x_v) = a_{\text{lcm}(\gamma_1, \dots, \gamma_m)}(x_v)$, we can see that the right-hand side above is equivalent to $x_v^{-m} \xi(x_v, \dots, x_v) f_1(1) \cdots f_m(1) f_{m+1}^{[\text{lcm}(\gamma_1, \dots, \gamma_m)]}(x_v)$. From formula (4.11), we thus complete the proof. $\square$

Example 4.9. Retaining the notation in Example 2.5, we have $S_{\mathbf{f}} = f \circ \text{gcd}$. Hence, by formula (4.12), we have for a factor-closed set $S = \{x_1, \dots, x_n\}$

$$\det_I((f \circ \text{gcd})(x_{i_1}, \dots, x_{i_{m+1}}))_{1 \leq i_1, \dots, i_{m+1} \leq n} = \prod_{v=1}^n (f * \mu)(x_v).$$

This is a part of the result of P. Haukkanen [10].

References

- [1] D. R. Anderson and T. M. Apostol, The evaluation of Ramanujan's sum and generalizations, *Duke Math. J.*, **20** (1953) 211–216.
- [2] G. E. Andrews, R. Askey and R. Roy, Special functions, Encyclopedia of Math. Appl., 71, Cambridge University Press, Cambridge, 1999.
- [3] T. M. Apostol, Arithmetical properties of generalized Ramanujan sums, *Pacific J. Math.* **41** (1972) 281–293.
- [4] J. M. Borwein, K. K. Choi, On Dirichlet Series for sums of squares, Rankin memorial issues, *Ramanujan J.*, **7** (2003) 95–127.

- [5] K. Bourque and S. Ligh, Matrices associated with classes of arithmetical functions, *J. Number Theory*, **45** (1993) 367–376.
- [6] D. Bump, Automorphic forms on $GL(3, \mathbb{R})$, Lecture Notes in Mathematics, 1083. Springer-Verlag, Berlin, 1984.
- [7] D. Bump, The Rankin-Selberg method: a survey, Number theory, trace formulas and discrete groups (Oslo, 1987), Academic Press, Boston, MA, 1989 49–109.
- [8] E. Cohen, A class of arithmetical functions, *Proc. Nat. Acad. Sci. U.S.A.*, **41** (1955) 939–944.
- [9] E. Cohen, A class of arithmetical functions in several variables with applications to congruences, *Trans. Amer. Math. Soc.*, **96** (1960) 335–381.
- [10] P. Haukkanen, Higher-dimensional GCD matrices, *Linear Algebra Appl.*, **170** (1992) 53–63.
- [11] K. Kimoto, K. Kurokawa, S. Matsumoto and M. Wakayama, Multiple finite Riemann zeta functions, *Acta Arith.* **116** (2005) 173–187.
- [12] P. J. McCarthy, A generalization of Smith’s determinant, *Canad. Math. Bull.*, **29** (1986) 109–113.
- [13] P. J. McCarthy, Introduction to arithmetical functions, Universitext. Springer-Verlag, New York, 1986.
- [14] K. Motose, Ramanujan’s sums and cyclotomic polynomials, *Math. J. Okayama Univ.*, **47** (2005) 65–74.
- [15] C. A. Nicol, Some formulas involving Ramanujan sums, *Canad. J. Math.*, **14** (1962) 284–286.
- [16] R. Oldenburger, Higher dimensional determinants, *Amer. Math. Monthly*, **47** (1940) 25–33.
- [17] S. Ramanujan, On certain trigonometrical sums and their applications in the theory of numbers, *Trans. Cambridge Phil. Soc.*, **22** (1918) 259–276.
- [18] L. H. Rice, Introduction to higher determinants, *J. Math. Phys.*, **9** (1930) 47–71.
- [19] H. J. S. Smith, On the value of a certain arithmetical determinant, *Proc. London Math. Soc.*, **7** (1875/76) 208–212.
- [20] E. C. Titchmarsh, The Theory of the Riemann Zeta-function, 2nd Ed., Oxford Science Publications, 1986.
- [21] R. Vaidyanathaswamy, The theory of multiplicative arithmetic functions, *Trans. Amer. Math. Soc.*, **33** (1931) 579–662.

YOSHINORI YAMASAKI

Graduate School of Mathematics, Kyushu University.

Hakozaki, Fukuoka, 812-8581 JAPAN.

yamasaki@math.kyushu-u.ac.jp